Opioid Meeting

07/26/2017

Medicaid Transformation Demonstration Toolkit Measures						Assessment of ACH Performance, by Demonstration Year		
Measure Name	**NQF#**	**Measure Steward**	**Measure Description**	**Method for Assessment of ACH Performance**	**Reporting Responsibility**	**DY 3 (2019)**	**DY 4 (2020)**	**DY 5 (2021)**
Inpatient Hospital Utilization		NCQA	For members 18 years of age and older, the risk-adjusted ratio of observed to expected acute inpatient discharges during the mesaurement year reported by Surgery, Medicine, and Total	Improvement over self	HCA	Inactive	P4P	P4P
Medication Assisted Therapy (MAT) with Buprenorphine or Methadone		Bree Collaborative	The count and percentage of Medicaid members with a documented diagnosis of opioid abuse/dependence who are engaged in MAT: Buprenorphine or Methadone	Improvement over self	HCA	P4P	P4P	P4P
Outpatient Emergency Department Visits per 1000 Member Months		NCQA / RDA	The rate of Medicaid enrollee visits to emergency department per 1000 member months, including visits related to mental health and chemical dependency. Separate reporting for age groups 10-17, 18-64, and 65+.	Improvement over self	HCA	P4P	P4P	P4P
Patients on high-dose chronic opioid therapy by varying thresholds (Measure specification in development)		Bree Collaborative	Measure specifications in development. Among Medicaid enrollees, percent of chronic opioid therapy patients receiving doses: >50 mg. MED in a quarter, doses >90 mg. MED in a quarter.	Improvement over self	HCA	P4P	P4P	P4P
Patients with concurrent sedatives prescriptions (Measure specification in development)		Bree Collaborative	Measure specifications in development. Among Medicaid enrollees receiving chronic opioid therapy, what percent had more than 45 days of Sedative Hypnotics/ Benzodiazepines/ carisoprodol/ barbiturates dispensed in the quarter.	Improvement over self	HCA	P4P	P4P	P4P
Substance Use Disorder Treatment Penetration (Opioid) (Measure specification in development)		RDA	Percent of Medicaid enrollees with a diagnosis of opioid use disorder who have a substance use service need who received at least one qualifying service during the measurement year. Reported separately for adults and for children.	Improvement over self	RDA	Inactive	P4P	P4P